

Great Lakes Gastroenterology

(440) 205-1225

Clenpiq Split Dose

It is IMPORTANT that you read these instructions well in advance and follow them completely to ensure a successful exam. It is critical for you to thoroughly clean out your colon. Stool remaining in the colon can hide polyps or other abnormalities.

- Fill your prescription for Clenpiq at least several days before your procedure. DO NOT refrigerate or freeze. Clenpiq is ready to drink eliminating the need to mix.

Due to regulatory requirements and your safety, you must arrange for a responsible adult to drive you home after the procedure. We will ask for contact information for your driver when you check in. Taxi cabs, public transportation and Uber do not meet this requirement unless you have a responsible person with you. Your procedure will be cancelled if you do not comply with this requirement.

	DAY BEFORE your procedure	DAY of your procedure
Diet Instructions	<ul style="list-style-type: none">• NO SOLID FOOD THE ENTIRE DAY• Drink clear liquids all day such as water, sport drinks, tea or black coffee (no milk or cream), broth, apple or white grape juice, sodas, jello, or popsicles. NO red or purple• Stay hydrated and drink plenty of fluids• NO ALCOHOL or any recreational drugs	<ul style="list-style-type: none">• You can drink clear liquids up to <u>4 hours</u> before your appointment.• No gum, candy or breath mints within <u>4 hours</u> of your appointment.• Do not use any form of tobacco products <u>4 hours</u> before your test.• Medications may be taken <u>4 hours</u> before your appointment with a small sip of water.• Remove contacts lenses, and all body piercings.
Bowel Preparation	<ul style="list-style-type: none">• Drink clear liquids throughout the day.• At 5pm the evening before, drink the entire first bottle of Clenpiq right from the bottle. Drinking the bottle slower will help prevent stomach upset.• Follow this dose by drinking at least five 8 ounce cups of clear liquid using the cup provided over the next five hours.	<ul style="list-style-type: none">• About 6 hours before your colonoscopy at _____ a.m. drink the entire second bottle of Clenpiq right from the bottle. Drinking the bottle slower will help prevent stomach upset.• Follow this dose by drinking at least three 8 ounce cups of clear liquid using the cup provided over the next two hours. You must stop drinking clear liquids _____ a.m. hours before your test.

-OVER-

Great Lakes Gastroenterology

(440) 205-1225

Medication Instructions:

Notify the office if you have an implanted heart defibrillator, pacemaker, or decreased kidney function. If you have diabetes and take insulin, contact your physician that manages this for specific instructions.

If you check your blood sugars, test it the evening before and the morning of your procedure.

HOLD the following medications using these guidelines, unless otherwise instructed by your Primary Care doctor, Cardiologist or Endocrinologist

	7 DAYS BEFORE	5 DAYS BEFORE	2 DAYS BEFORE
Blood Thinners		<ul style="list-style-type: none">• Warfarin (Coumadin)• Anti-platelets such as Clopidogrel/Plavix, Ticlid, Effient, Brilinta, Aggrenox, Pletal, Arixtra, Elmiron, Persantine Ticlopidine	<ul style="list-style-type: none">• Pradaxa, Xarelto, Eliquis, Savaysa
Iron	<ul style="list-style-type: none">• Iron supplements (ferrous sulfate)		

Medications to **HOLD** the morning of the procedure

Oral Diabetes Meds	<ul style="list-style-type: none">• Metformin, glipizide, glucophage, glucotrol, glyburide, tolazamide, tolbutamide
Blood Pressure & Heart Meds	<ul style="list-style-type: none">• Ace Inhibitors: ending in 'pril' (Lisinopril, Ramipril, Enalapril or Vasotec, Moexipril or Univas, Captopril or Capoten, Perindopril or Aceon, Accupril)• Angiotensin II Receptor Blockers (ARB's): ending in 'artan' (Losartan, Valsartan, Benicar)• Angiotensin II Receptor Blockers (ARB 's) combined with calcium channel blockers such as (Amlodipine/Valsartan, Amlodipine/Olmesartan, and Telmisartan/Amlodipine)
Diuretics Water pills	<ul style="list-style-type: none">• HCTZ (hydrochloro-thiazide), Lasix (furosemide), Maxzide, Chlorthalidone, Dyazide (triamterene/hydrochlorothiazide), Spironolactone

****TAKE ALL OTHER BLOOD PRESSURE MEDICATIONS NOT LISTED ABOVE****

- If you use a rescue inhaler, please be sure to bring it with you
- Bring your current insurance card(s), photo ID, and any copay that is due
- The Endoscopy Center opens at 7am. Please do not arrive any earlier than 7am

Procedure Date: ____/____/____

Approximate Arrival Time: ____:____ a.m / p.m.

Special Instructions: _____

You will receive a call the week day prior to your procedure with the actual arrival time

Tri Point - Main Entrance
7590 Auburn Rd.
Concord 44077

Endoscopy Center of Lake County, LLC
9614 Old Johnnycake Ridge Rd.
Mentor 44060

Lake West - Same Day Surgery
36000 Euclid Avenue
Willoughby 44094