

Great Lakes Gastroenterology

(440) 205-1225

Nu-Lytely

It is IMPORTANT that you read these instructions well in advance and follow them completely to ensure a successful exam. It is critical for you to thoroughly clean out your colon. Stool remaining in the colon can hide polyps or other abnormalities.

- Obtain in advance: Nu-lytely prescription from your pharmacy

Due to regulatory requirements and your safety, you must arrange for a responsible adult to drive you home after the procedure, We will ask for contact information for your driver when you check in. Taxi cabs, public transportation and Uber do not meet this requirement unless you have a responsible person with you. Your procedure will be cancelled if you do not comply with this requirement.

	DAY BEFORE your procedure	DAY of your procedure
Diet Instructions	<ul style="list-style-type: none"> • NO SOLID FOOD THE ENTIRE DAY • NO ALCOHOL or any recreational drugs • Drink clear liquids all day such as water, sport drinks, tea or black coffee (no milk or cream), broth, apple or white grape juice, soda¹s, jello, or popsicles. NO red or purple 	<ul style="list-style-type: none"> • You can drink clear liquids up to <u>4 hours</u> before your appointment. • No gum, candy or breath mints within <u>4 hours</u> of your appointment. • Do not use any form of tobacco products <u>4 hours</u> before your test • Medications may be taken <u>4 hours</u> before your appointment. • Remove contacts lenses, and all body piercings.
Bowel Preparation	<p style="text-align: center;"><u>Procedure scheduled before 1:00 pm:</u></p> <ul style="list-style-type: none"> • Drink clear liquids throughout the day. • In the morning, mix Nu-lytely and refrigerate. • At 4:00pm start drinking the bowel prep, one 8 oz. glass every 10-15 minutes until gone. 	
	<p style="text-align: center;"><u>Procedure scheduled at or after 1:00 pm</u></p> <ul style="list-style-type: none"> • Drink clear liquids throughout the day. • In the morning, mix Nu-lytely and refrigerate. • At 4:00pm start drinking the bowel prep, one 8 oz. glass every 10-15 minutes until 3/4 of the container is gone. This will take 4 hours. 	<p style="text-align: center;"><u>Procedure scheduled at or after 1:00 pm</u></p> <ul style="list-style-type: none"> • At _____ (5 hours prior to your arrival time), drink the remaining 1/4 of the prep, one 8 oz. glass every 10-15 minutes. Should be gone in 1 hour.

Great Lakes Gastroenterology

(440) 205-1225

Medication Instructions:

Notify the office if you have an implanted heart defibrillator, pacemaker, or decreased kidney function. If you have diabetes and take insulin, contact your physician that manages this for specific instructions.

If you check your blood sugars, test it the evening before and the morning of your procedure.

HOLD the following medications using these guidelines, unless otherwise instructed by your Primary Care doctor, Cardiologist or Endocrinologist

	7 DAYS BEFORE	5 DAYS BEFORE	2 DAYS BEFORE
Blood Thinners		<ul style="list-style-type: none"> Warfarin (Coumadin) Anti-platelets such as Clopidogrel/Plavix, Ticlid, Effient, Brilinta, Aggrenox, Pletal, Arixtra, Elmiron, Persantine Ticlopidine 	<ul style="list-style-type: none"> Pradaxa, Xarelto, Eliquis, Savaysa
Iron	<ul style="list-style-type: none"> Iron supplements (ferrous sulfate) 		

Medications to **HOLD the morning** of the procedure

Oral Diabetes Meds	<ul style="list-style-type: none"> Metformin, glipizide, glucophage, glucotrol, glyburide, tolazamide, tolbutamide
Blood Pressure & Heart Meds	<ul style="list-style-type: none"> Ace Inhibitors: ending in 'pril' (Lisinopril, Ramipril, Enalapril or Vasotec, Moexipril or Univasc, Captopril or Capoten, Perindopril or Aceon, Accupril) Angiotensin II Receptor Blockers (ARB's): ending in 'artan' (Losartan, Valsartan, Benicar) Angiotensin II Receptor Blockers (ARB's) combined with calcium channel blockers such as (Amlodipine/Valsartan, Amlodipine/Olmesartan, and Telmisartan/Amlodipine)
Diuretics Water pills	<ul style="list-style-type: none"> HCTZ (hydrochloro-thiazide), Lasix (furosemide), Maxzide, Chlorthalidone, Dyazide (triamterene/hydrochlorothiazide), Spironolactone

****TAKE ALL OTHER BLOOD PRESSURE MEDICATIONS NOT LISTED ABOVE****

- If you use a rescue inhaler, please be sure to bring it with you
- Bring your current insurance card(s), photo ID, and any copay that is due
- The Endoscopy Center opens at 7am. Please do not arrive any earlier than 7am

Procedure Date: ____/____/____

Approximate Arrival Time: ____:____ a.m / p.m.

Special Instructions: _____

You will receive a call the week day prior to your procedure with the actual arrival time

Tri Point - Main Entrance
7590 Auburn Rd.
Concord 44077

Endoscopy Center of Lake County, LLC
9614 Old Johnnycake Ridge Rd.
Mentor 44060

Lake West - Same Day Surgery
36000 Euclid Avenue
Willoughby 44094